

You must fill out the information below to receive your results and possible CME credit. You must correctly answer 70% or more of the five questions below to be awarded CME credit. Submit completed test to [jenny@societyhq.com](mailto:jenny@societyhq.com) for scoring.

Name:

AOA #:

Email Address:

Institution:

### **Module 8: Osteopathic Pediatrics**

1. Use of which of the following OMM techniques has been shown to result in increased sIgA levels?
  - A. CV-4
  - B. HVLA
  - C. Muscle energy
  - D. Occipitoatlantal release
  - E. Strain counterstrain
  
2. Which of the following is an expected finding during the osteopathic exam of a 12 month old infant's head?
  - A. Asymmetry of facial features
  - B. Closed anterior fontanelle
  - C. Motion of individual cranial bones
  - D. Open posterior fontanelle
  - E. Overriding sutures
  
3. At what age can segmental somatic dysfunction be diagnosed in a child?
  - A. 6 days old
  - B. 6 weeks old
  - C. 6 months old
  - D. 6 years old
  - E. 16 years old

4. Cranial OMM has been found to be efficacious in the treatment of which of the following pediatric conditions?
  - A. Colic
  - B. Concussion
  - C. Depression
  - D. Global developmental delay
  - E. Migraine
  
5. A 4 year old child presents with musculoskeletal rib pain after an injury. Which of the following OMM techniques is most likely beneficial without the possibility of an adverse outcome?
  - A. CV-4 cranial dysfunction
  - B. HVLA cervical dysfunction
  - C. Lymphatic pump
  - D. Muscle energy lumbar dysfunction
  - E. Myofascial release thoracic dysfunction